

What is claimed is:

1. An isolated nucleic acid molecule selected from the group consisting of:

- 5 a) a nucleic acid molecule comprising a nucleotide sequence which is at least 60% identical to the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, or a complement thereof;
- 10 b) a nucleic acid molecule comprising a fragment of at least 583 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, or a
- 25

30 c) a polypeptide molecule comprising an amino acid sequence at least about 60% identical to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID

- 150 -

NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316;

d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ

98991, 98993, 98994, or PTA-316; and the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316; and

- 151 -

e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, or the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, under stringent conditions.

- 152 -

2. The isolated nucleic acid molecule of claim 1 which is selected from the group consisting of:

5 a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, or the DNA
10 insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, or a complement thereof; and

15 b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ
20 ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316.

25

3. The nucleic acid molecule of claim 1 further comprising vector nucleic

acid sequence

4. The nucleic acid molecule of claim 1 further comprising nucleic acid
30 sequences encoding a heterologous polypeptide.

- 153 -

5. A host cell which contains the nucleic acid molecule of claim 1.

6. The host cell of claim 5 which is a mammalian host cell.

5 7. A non-human mammalian host cell containing the nucleic acid molecule of claim 1.

8. An isolated polypeptide selected from the group consisting of:

10 a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72.

15 20 25 30

PLASMID DEPOSITED WITH ATCC AS ACCESSION NUMBER 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316;

- 154 -

b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, or the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316 under stringent conditions; and

c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, or the DNA insert of the plasmid deposited

- 155 -

with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316.

5 d) a polypeptide comprising an amino acid sequence which is at least 60% identical to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, 10 SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316.

15

9. The isolated polypeptide of claim 8 comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 20 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316.

25

30 11. An antibody which selectively binds to a polypeptide of claim 8.

12. A method for producing a polypeptide selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316;
- b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the

- 157 -

plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316; and

- c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:70, or SEQ ID NO:72 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:58, SEQ ID NO:69, or SEQ ID NO:71, or the DNA insert of the plasmid deposited with ATCC as Accession Number 98936, 98937, 98938, 98939, 98940, 98941, 98942, 98943, 98944, 98945, 98946, 98947, 98948, 98949, 98950, 98951, 98991, 98993, 98994, or PTA-316 under stringent conditions;

- 158 -

13. A method for detecting the presence of a polypeptide of claim 8 in a sample comprising:

- a) contacting the sample with a compound which selectively binds to the polypeptide; and
- 5 b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 8 in the sample.

14. The method of claim 13, wherein the compound which binds to the
10 polypeptide is an antibody.

15. A kit comprising a compound which selectively binds to a polypeptide of claim 8 and instructions for use.

15 16. A method for detecting the presence of a nucleic acid molecule in claim 1 in a sample comprising:

- a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and
- 20 b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of claim 1 in the sample.

17. The method of claim 16, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

25

18. A kit comprising a compound which selectively hybridizes to a nucleic

19. A method for identifying a compound which binds to a polypeptide of claim 8 comprising:

- a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and
- 5 b) determining whether the polypeptide binds to the test compound.

20. The method of claim 19, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detection of test compound/polypeptide binding;
- 10 b) detection of binding using a competition binding assay; and
- c) detection of binding using an assay for PCIP activity.

21. A method for modulating the activity of a polypeptide of claim 8 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

22. A method for identifying a compound which modulates the activity of a polypeptide of claim 8 comprising:

- a) contacting a polypeptide of claim 8 with a test compound; and
- 20 b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

23. A method for identifying a compound capable of treating a disorder

comprising assaying a polypeptide or a cell expressing the polypeptide with the PCIP nucleic acid molecule of claim 1 for the activity of the PCIP polypeptide of claim 8, thereby identifying a compound capable of treating a disorder characterized by aberrant PCIP nucleic acid expression or PCIP protein activity.

24. The method of claim 23, wherein the disorder is a CNS disorder.
25. The method of claim 24, wherein the disorder is epilepsy.
27. The method of claim 24, wherein the disorder is spinocerebellar ataxia.
28. The method of claim 23, wherein the disorder is a cardiovascular disorder.
29. The method of claim 28, wherein the cardiovascular disorder is associated with an abnormal I_{to} current.
30. A method for determining if a subject is at risk for a disorder characterized by aberrant or abnormal PCIP nucleic acid expression and/or PCIP protein activity comprising detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion, wherein the genetic lesion is characterized by an alteration affecting the integrity of a gene encoding the PCIP polypeptide of claim 8 or misexpression of the PCIP nucleic acid molecule of claim 1.
31. The method of claim 30, wherein the disorder is a CNS disorder.
32. The method of claim 31, wherein the disorder is epilepsy.
33. The method of claim 31, wherein the disorder is spinocerebellar ataxia.
35. The method of claim 34, wherein the cardiovascular disorder is associated with an abnormal I_{to} current.

- 161 -

36. A method for identifying a subject suffering from a disorder characterized by aberrant or abnormal PCIP nucleic acid expression and/or PCIP protein activity comprising obtaining a biological sample from the subject, and detecting in the sample, the presence or absence of a genetic lesion, wherein the genetic lesion is characterized by an alteration affecting the integrity of a gene encoding the PCIP polypeptide of claim 8 or misexpression of the PCIP nucleic acid molecule of claim 1, thereby identifying a subject suffering from a disorder characterized by aberrant or abnormal PCIP nucleic acid expression and/or PCIP protein activity.
37. The method of claim 36, wherein the disorder is a CNS disorder.
38. The method of claim 37, wherein the disorder is epilepsy.
39. The method of claim 37, wherein the disorder is spinocerebellar ataxia.
40. The method of claim 36, wherein the disorder is a cardiovascular disorder.
41. The method of claim 40, wherein the cardiovascular disorder is associated with an abnormal I_{to} current.
42. A method for treating a subject having a potassium channel associated disorder comprising administering to the subject a PCIP polypeptide of claim 8 or portion thereof such that treatment occurs.
44. The method of claim 43, wherein the disorder is epilepsy.
45. The method of claim 43, wherein the disorder is spinocerebellar ataxia.

46. The method of claim 42, wherein the disorder is a cardiovascular disorder.

5 47. The method of claim 46, wherein the cardiovascular disorder is associated with an abnormal I_{to} current.

48. A method for treating a subject having a potassium channel associated disorder comprising administering to the subject a nucleic acid encoding a PCIP
10 polypeptide of claim 8 or portion thereof such that treatment occurs.

49. The method of claim 48, wherein the disorder is a CNS disorder.

50. The method of claim 49, wherein the disorder is epilepsy.

15

51. The method of claim 49, wherein the disorder is spinocerebellar ataxia.

52. The method of claim 48, wherein the disorder is a cardiovascular disorder.

20

53. The method of claim 52, wherein the cardiovascular disorder is associated with an abnormal I_{to} current.

54. Use of the compound identified in the method of claim 23 to treat a
25 potassium channel associated disorder.